



1/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form A measured with variable slits.

Fig. 1


2/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form B measured with variable slits.

Fig. 2


3/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form C measured with variable slits.

Fig. 3


4/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form E measured

Fig. 4


5/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form F measured with variable slits.

Fig. 5


6/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form G measured with variable slits.

Fig. 6


7/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form H measured with variable slits.

Fig. 7


8/9



A Raman spectrum of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form A

Fig. 8

9/9



A Raman spectrum of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazol[1,2-a]pyridine-6-carboxamide mesylate salt form B

Fig. 9